



The yellow yolk in the egg provides the chick with food.

Birds and reptiles often have a temporary horn-like 'tooth' to help them break out of the egg.

A clutch is the name for a number of eggs that a bird lays in one go.

The ostrich egg is the biggest in the world.

The smallest egg in the world is laid by the bee hummingbird.

- The great spotted woodpecker incubates its egg for only ten days.
- The first fossilized dinosaur eggs found were 1 foot (30 cm) long, 10 inches (25 cm) wide, had a volume of about half a gallon (2 litres), and may have weighed up to 15.5 pounds (7 kg).
- When animals started to walk on land, the soft-shelled eggs of amphibians became hard shelled eggs of reptiles for increased protection and to stop the egg drying out.



The yellow yolk in the egg provides the chick with food.

Birds and reptiles often have a temporary horn-like 'tooth' to help them break out of the egg.

A clutch is the name for a number of eggs that a bird lays in one go.

The ostrich egg is the biggest in the world.

The smallest egg in the world is laid by the bee hummingbird.

- The great spotted woodpecker incubates its egg for only ten days.
- The first fossilized dinosaur eggs found were 1 foot (30 cm) long, 10 inches (25 cm) wide, had a volume of about half a gallon (2 litres), and may have weighed up to 15.5 pounds (7 kg).
- When animals started to walk on land, the soft-shelled eggs of amphibians became hard shelled eggs of reptiles for increased protection and to stop the egg drying out.